

OPERATIONAL QUANTUM GRAVITY FOR ENGINEERS

A revised damping, vacuum-polarizability, and uncertainty-based interpretation of gravitational scaling

Todd J. Desiato

Statesville, North Carolina, USA

Revised and consolidated from earlier manuscripts and supporting literature

V11_final

April 11, 2026

Abstract

Practically speaking, time is what clocks measure and space is what rulers measure. In this paper I restate and consolidate my long-running program in the strongest form I believe can be defended rigorously from my manuscript series and the primary literature on which it builds. My central claim is interpretive rather than iconoclastic: wherever the same field relations and scaling laws are recovered, the model makes the same observable predictions as general relativity, but it assigns a different physical meaning to those relations. In the present reading, the metric is a compact mathematical encoding of comparisons among physical clocks, rulers, signals, frequencies, and energies. It need not be taken as proof that a spacetime manifold is the unique fundamental ontology. I summarize the static weak-field transformations of the Schwarzschild / polarizable-vacuum representation and show that the adopted scaling preserves the Heisenberg products $\Delta x \Delta p$ and $\Delta t \Delta E$ exactly. I then write my radiative-damping model in a form that reproduces the same transformation table through the identification $K=(1-\zeta^2)^{-1}$. The controlling quantity is treated as a real effective scalar order parameter built from the local stochastic electromagnetic-magnetic environment seen by matter, and I add a phenomenological weak-field source law together with a linear-response bridge from the spectral environment $S_{\text{env}}(\omega, x)$ to the damping variable $\zeta(x)$. The paper also states explicitly how universality of free fall can emerge at leading order when the K -map is common to all matter processes, and it sharpens the experimental program in precision spectroscopy, clock comparison, and resonance-based perturbation studies. Thermodynamic analogies are used as motivation, not as proof. The resulting synthesis is not claimed as a completed microscopic theory of gravitation. It is offered as a mathematically coherent operational interpretation of the same observed weak/static relations, together with a more explicit engineering research program for testing whether a deeper non-geometrical layer exists.

Keywords: operational time, quantum gravity, polarizable vacuum, variable refractive index, stochastic vacuum environment, radiative damping, uncertainty principle, weak-field reinterpretation, precision metrology

Nomenclature

Symbol	Meaning	Remarks
$K(x)$	effective polarizable-vacuum / metric scaling parameter	adopted static coordinate control variable
$\zeta(x)$	relative damping factor	phenomenological microscopic matching variable
c_0	local speed of light in an unperturbed local inertial frame	taken invariant locally
c_K	coordinate speed of light as compared by a distant observer	equals c_0/K in the adopted convention
$\Delta x, \Delta t$	coordinate-comparison length and time increments	not ontological primitives
$\Delta p, \Delta E$	momentum and energy uncertainties	chosen to preserve Heisenberg products
$S_{\text{env}}(\omega, x)$	effective local spectral environment seen by matter	baseline ZPF plus matter-generated stochastic loading
m_{eff}	effective coordinate mass parameter	not a claim that local invariant rest mass changes
$\Phi_{\zeta}(x)$	weak-field potential associated with $\zeta(x)$	defined by $\zeta^2 = -2\Phi_{\zeta}/c_0^2$ in the weak/static closure
$\rho_{\text{eff}}(x)$	effective source density for the weak-field closure	matter density plus matter-induced environmental loading
$u_{\text{env}}(x)$	coarse-grained local environmental energy density	weighted integral over the effective spectral environment
$W(\omega), G_{\gamma}(\omega)$	spectral weighting and response kernels	phenomenological functions connecting S_{env} to γ_{eff} and ζ
η_{ab}	Eötvös parameter for compositions a and b	used to state universality-of-free-fall bounds

1. Introduction

Practically speaking, a clock compares rates and a ruler compares lengths. The quantities that enter gravitation are therefore operational comparisons among physical processes, not substances called time and space. General relativity encodes those comparisons in a metric, and I do not dispute its empirical success. My question is narrower: can the same observational content be written in a more direct engineering language based on matter, vacuum response, damping, and scale-setting processes? [3-5,19-23]

My manuscript series from 2006 through 2023 follows one line of development. First, the weak/static transformations associated with the Schwarzschild solution can be expressed through a single scalar quantity K in a polarizable-vacuum representation. Second, the same table of transformations can be written so that the Heisenberg products remain invariant. Third, the same table can be matched again by introducing a damping factor ζ for matter treated as a driven oscillator in stochastic equilibrium with its environment. The strongest version of the program is therefore not that geometry is wrong, but that geometry may be descriptive rather than ontologically unique. [1,2,4,5,19-23]

That distinction matters. If the metric is descriptive rather than ontologically unique, then a deeper engineering model should begin with observables and then ask what microscopic changes in matter would cause the observed changes in clock rates, characteristic frequencies, lengths, and energies. My interpretive move is operational: time is what clocks measure; length is what rulers measure; the metric is a concise bookkeeping device for those relations; and a deeper causal layer may be sought in how matter is driven, damped, and scaled by its local environment. [1,2,4,5,12,19-23]

Stated plainly, I am not arguing that general relativity is empirically wrong in the regime treated here. Wherever the same field relations and scaling laws are recovered, the observable content of my model is identical to that of GR. What changes is the ontology attributed to the equations. As an interpretation of the same successful weak/static relations, I regard the present model as standing on equal empirical footing with the standard geometrical reading so long as both recover the same observational table. [3-5,19-23]

On this reading, Einstein's field equation may be approached operationally before it is approached ontologically. The geometric side $G_{\mu\nu}$ summarizes the relational structure inferred from clock comparisons, ruler comparisons, signal propagation, and material motion. The matter side $T_{\mu\nu}$ describes the stress-energy content of the matter fields from which those same clocks, rulers, and signals are physically built. In that sense the equation need not be read first as a declaration that a spacetime manifold is the fundamental substance of nature. It may instead be read as a comparison law linking observed relational structure to the matter sector that sets the standards of observation itself. [3-5,12,19-23]

I do not regard the geometrical interpretation as a mistake in any simple sense. It has enriched the mathematics of gravitation enormously. My narrower concern is that its ontological primacy may also have constrained the search for quantum gravity by encouraging us to treat geometry as fundamental rather than emergent. One advantage of the present interpretation is economy: it keeps the successful mathematics where it works, but asks first what clocks do, what rulers do, and how quantum matter sets their scales. [3-5,12,19-23]

2. Operational starting point and the metric / PV correspondence

The polarizable-vacuum line of thought runs from Wilson and Dicke to Puthoff's variable- K representation of static gravitational effects. In that representation the vacuum is treated as an effective medium whose single scalar parameter K summarizes how clocks, rods, and the coordinate speed of light compare between an altered region and a distant unaltered one. For my purposes this is attractive because it preserves the

observational content while replacing purely geometrical language with a form more natural to engineering analysis. [1,2,4-6]

In this paper I adopt the same static line element in the spirit of my earlier papers. I do not present it as a new derivation of general relativity; I present it as the operational encoding of the same weak/static gravitational scaling relations that I wish to reinterpret microscopically. [4,5,19,22,23]

$$ds^2 = \frac{c_0^2 dt^2}{K(x)} - K(x)(dx^2 + dy^2 + dz^2) \quad (1)$$

$$\frac{c_K}{c_0} = \frac{1}{K}, \quad \frac{\Delta x_K}{\Delta x_0} = \frac{1}{\sqrt{K}}, \quad \frac{\Delta t_K}{\Delta t_0} = \sqrt{K}, \quad \frac{\omega_K}{\omega_0} = \frac{1}{\sqrt{K}} \quad (2)$$




Figure 1. Operational equivalence ladder for the interpretive model.

The point of the ladder in Fig. 1 is interpretive rather than decorative. The macroscopic metric coefficients, the PV refractive-index variable K , and the microscopic damping picture are treated here as different encodings of the same observed weak/static comparison rules. In that restricted sense the present model is a reinterpretation of the GR table, not a competing table. [4,5,22,23]

3. Uncertainty-compatible scaling

Once I adopt the operational statements that lengths are contracted and clocks are slowed in the chosen coordinate comparison, the uncertainty products must still be respected. This is the key point. The uncertainty principle constrains products, not isolated variables. The mathematical question is therefore whether one can choose the complementary scalings of momentum and energy so that the products $\Delta x \Delta p$ and $\Delta t \Delta E$ remain unchanged while the observable table of gravitational scalings is reproduced. [14-17,21-23]

One can. The proof is immediate by multiplication. This does not, by itself, derive the full gravitational field equations. It does something more modest and more secure: it shows that the scaling map I adopt is

compatible with the kinematical quantum constraints. In that sense the uncertainty principle enters here as a consistency condition on the scaling table rather than as a stand-alone first-principles derivation of gravity. [14-17,21]

An important consequence follows. If one further demands that force remain invariant in the adopted comparison scheme, then the quantity that scales like mass cannot be treated naively. What emerges is an effective coordinate mass parameter. I keep that point explicit to avoid the common misunderstanding that I am claiming the local invariant rest mass of a particle changes in its own local inertial frame. My claim is weaker and cleaner: within the distant-observer bookkeeping used here, an effective mass parameter must scale as $K^{3/2}$ if the rest of the table is to remain internally consistent. [21-23]

$$\Delta x_K \Delta p_K = \Delta x_0 \Delta p_0, \quad \Delta t_K \Delta E_K = \Delta t_0 \Delta E_0 \quad (3)$$

$$\Delta p_K = \sqrt{K} \Delta p_0, \quad \Delta E_K = \Delta E_0 / \sqrt{K} \quad (4)$$

$$\frac{v_K}{v_0} = \frac{1}{K}, \quad \frac{P_K}{P_0} = \frac{1}{K}, \quad \frac{a_K}{a_0} = \frac{1}{K^{3/2}}, \quad \frac{m_{\text{eff},K}}{m_0} = K^{3/2} \quad (5)$$

4. Damped oscillator model and matter scale

In my later manuscripts I move from a purely kinematic table to a microscopic picture in which matter is treated, for engineering purposes, as an ensemble of oscillators. In that picture the quantum vacuum supplies a baseline driving field while radiative damping and local environmental loading alter the steady-state equilibrium. This move is motivated by the literature on vacuum-fluctuation physics and by equilibrium accounts of radiationless quantum ground states. It is also motivated by the fact that oscillators provide a natural language for frequency, linewidth, power flow, and resonance. [8-10,20,22,23]

The central equation is the driven damped oscillator. From it I introduce the dimensionless damping factor $\zeta = \gamma/\omega_0$ and the underdamped frequency $\omega_\zeta = \omega_0 \sqrt{1-\zeta^2}$. In my engineering model the reduction in available driving power and the shift in characteristic energy are then mapped onto the same table of gravitational observables. The model does not merely borrow the standard undamped harmonic-oscillator ground state. It introduces a phenomenological fluctuation map in which the mean-square position, velocity, and acceleration fluctuations scale with powers of $(1-\zeta^2)$. That is the step that allows the damping picture to reproduce the same operational relations that K already encodes. [8-10,22,23]

The identification $K = (1-\zeta^2)^{-1}$ is therefore the hinge of the synthesis. Once that substitution is made, the frequency, energy, velocity, acceleration, and effective-mass entries of the damping table collapse onto the same operational relations as the metric / PV table. That is the strongest mathematical equivalence in the framework. What remains open is the microscopic origin of ζ and the exact field theory behind it. [22,23]

It is also useful to say explicitly that the stochastic and spectral pictures of the vacuum are not competing ontologies. A stochastic field can always be described by its spectrum, and a weighted spectral density is one natural coarse-graining of a random background. My present damping language therefore extends, rather than rejects, the earlier vacuum-equilibrium language in which particle scale was associated with equilibrium against a structured vacuum spectrum. [8-10,22-24]

In the older EGM harmonic work, the harmonic cut-off was interpreted as indicating the energy density at which matter reaches equilibrium with the surrounding polarizable vacuum / zero-point field. I do not

present that as a proof of the current model. I cite it because it captures the same physical intuition that motivates the present one: matter scale is not arbitrary, but is set by its equilibrium with a structured vacuum environment. [24]

$$m\ddot{x} + 2m\gamma\dot{x} + m\omega_0^2x = F_{\text{env}}(t) \quad (6)$$

$$\zeta \equiv \gamma/\omega_0, \quad \omega_\zeta = \omega_0\sqrt{1-\zeta^2}, \quad 0 \leq \zeta < 1 \quad (7)$$

$$E_\zeta = E_0\sqrt{1-\zeta^2}, \quad P_\zeta = P_0(1-\zeta^2) \quad (8)$$

$$\langle x^2 \rangle_\zeta = \langle x^2 \rangle_0(1-\zeta^2), \quad \langle v^2 \rangle_\zeta = \langle v^2 \rangle_0(1-\zeta^2)^2, \quad \langle a^2 \rangle_\zeta = \langle a^2 \rangle_0(1-\zeta^2)^3 \quad (9)$$

$$K(x) = \frac{1}{1-\zeta(x)^2} \quad (10)$$




Figure 2. Damping-parameter scalings used in the engineering model.

Table 1. Unified scaling map used throughout the paper

Quantity	Metric / PV row	Uncertainty-compatible row	Damping row
Speed of light	$c_K / c_0 = 1/K$	derived from adopted operational convention	$c_\zeta / c_0 = \sqrt{1-\zeta^2}$
Length	$\Delta x_K / \Delta x_0 = 1/\sqrt{K}$	chosen observable input	$\Delta x_\zeta / \Delta x_0 = \sqrt{1-\zeta^2}$
Time	$\Delta t_K / \Delta t_0 = \sqrt{K}$	chosen observable input	$\Delta t_\zeta / \Delta t_0 = 1/\sqrt{1-\zeta^2}$
Momentum	$\Delta p_K / \Delta p_0 = \sqrt{K}$	preserves $\Delta x \Delta p$	$\Delta p_\zeta / \Delta p_0 = 1/\sqrt{1-\zeta^2}$
Energy / frequency	$\Delta E_K / \Delta E_0 = \omega_K / \omega_0 = 1/\sqrt{K}$	preserves $\Delta t \Delta E$	$\Delta E_\zeta / \Delta E_0 = \omega_\zeta / \omega_0 = \sqrt{1-\zeta^2}$
Velocity / power	$v_K / v_0 = P_K / P_0 = 1/K$	ratio of length to time; energy to time	$v_\zeta / v_0 = P_\zeta / P_0 = 1-\zeta^2$
Acceleration	$a_K / a_0 = 1/K^{3/2}$	from F invariant and m_{eff} row	$a_\zeta / a_0 = (1-\zeta^2)^{3/2}$
Effective coordinate mass	$m_{\text{eff},K} / m_0 = K^{3/2}$	bookkeeping variable, not local rest mass	$m_{\text{eff},\zeta} / m_0 = (1-\zeta^2)^{-3/2}$

5. Effective stochastic background field and scalar control parameter

In earlier drafts I used the phrase “Maxwell Temporal Field” as a heuristic label. Here I make the intended meaning explicit. The controlling quantity is treated as a real effective scalar order parameter, or coarse-grained state variable, built from the local stochastic electromagnetic and magnetic environment seen by matter. This is the formulation most faithful to what I am actually proposing. [10,11,20,22,23]

I do not deny the baseline zero-point field of QED. My claim is that the actual environment experienced by matter is not an idealized empty baseline; it is loaded by surrounding matter, radiation, and internal hadronic and electronic activity. The local state seen by an atom or nucleus is therefore a real spectral environment, not a mere notational convenience. In this paper I represent the control variable as a functional of that environment. [8-10,20,22,23]

I retain the language of scalar magnetic flux only as engineering shorthand for the magnetic sector of that stochastic environment. What matters here is the weaker and more defensible statement: matter couples to a local stochastic field environment with a scalar control parameter capable, in principle, of shifting equilibrium frequency, available power, and fluctuation scale. That statement is sufficient for the present synthesis. [11,20,22,23]

It is equally important to say that the local stochastic vacuum and the local spectral vacuum are, in this framework, two ways of describing the same underlying physical layer. The stochastic description emphasizes loading, fluctuation amplitude, and local field variability. The spectral description emphasizes how that same environment is distributed in frequency. The present paper uses the stochastic language more often because it is better suited to damping, linewidth, response, and control; but I do not regard the two descriptions as physically separate. [10,20,22-24]

5.1 From spectral environment to damping

The key point is that the stochastic background is not the same thing as the damping variable. The local spectrum is the environment; ζ is the coarse-grained dissipative response of matter to that environment. Near equilibrium it is natural to summarize the environment by a weighted spectral density and to summarize the response by an effective damping rate. I therefore introduce the auxiliary quantities $u_{\text{env}}(\mathbf{x})$ and $\gamma_{\text{eff}}(\mathbf{x})$ as phenomenological state variables, not as a completed microscopic derivation. [8-10,20,22-25]

$$u_{\text{env}}(\mathbf{x}) = \int_0^\infty W(\omega) S_{\text{env}}(\omega, \mathbf{x}) d\omega, \quad (11)$$

$$\gamma_{\text{eff}}(\mathbf{x}) = \gamma_0 + \int_0^\infty G_\gamma(\omega) \delta S_{\text{env}}(\omega, \mathbf{x}) d\omega$$

$$\zeta(\mathbf{x}) = \gamma_{\text{eff}}(\mathbf{x})/\omega_0 \quad (12)$$

Here $W(\omega)$ and $G_\gamma(\omega)$ are weighting and response kernels that encode which portion of the local spectral environment is relevant to the mode under consideration. This formulation makes explicit what is only implicit in my earlier papers: the environment sets both fluctuation amplitude and dissipative response, while ζ is the scalar summary that enters the operational K-map. [8-10,20,22,23,25]

5.2 Minimal weak-field source law

In the absence of a completed microscopic field theory, I adopt a phenomenological weak-field closure. The minimal requirement is that the source law reduce to the ordinary Newtonian potential for static spherical matter while still allowing the environmental language of the present model. I therefore define a weak-field potential $\Phi_\zeta(\mathbf{x})$ by

$$\zeta^2(\mathbf{x}) = -2\Phi_\zeta(\mathbf{x})/c_0^2, \quad \nabla^2 \Phi_\zeta(\mathbf{x}) = 4\pi G \rho_{\text{eff}}(\mathbf{x}) \quad (13)$$

and take the effective source to be

$$\begin{aligned} \rho_{\text{eff}}(\mathbf{x}) &= \rho_m(\mathbf{x}) + \lambda \delta u_{\text{env}}(\mathbf{x})/c_0^2, \\ g(\mathbf{x}) &= -\nabla \Phi_\zeta(\mathbf{x}) = (c_0^2/2) \nabla \zeta^2(\mathbf{x}) \end{aligned} \quad (14)$$

These equations are not yet a finished microscopic theory. They are the minimal closure that connects matter distribution to the scalar control variable in the weak/static limit. For a point or spherically symmetric source, they reduce to the same leading-order dependence used later in Eq. (19). The baseline ZPF drops out of the source term; what matters is the local departure from the unperturbed environment together with the matter distribution that generates it. [3-5,20,22-25]

5.3 Radiative equilibrium, radiation reaction, and the emergence of free fall

Here I want to state more clearly how the equivalence principle enters the model. In the unperturbed state, matter is treated as residing in local radiative equilibrium with its environment. The equilibrium scale is set by a stationarity condition on the full matter-plus-environment system: mean input power from the local field environment is balanced by mean radiative and dissipative output power of the bound system. In the coarse-grained language adopted here, that balance may be written as

$$P_{\text{in}}(x) = P_{\text{out}}(x), \quad \delta P(x) \equiv P_{\text{in}}(x) - P_{\text{out}}(x) = 0 \quad (15)$$

I regard this local stationarity, not geometry by itself, as the symmetry relevant to scale setting. Stated carefully, the corresponding Noether statement belongs to the underlying closed matter-plus-environment system rather than to the reduced damped subsystem alone. The present paper therefore does not claim a formal Noether derivation at the coarse-grained level. It claims the weaker point that free fall can be interpreted as a gradient-driven departure from a locally stationary equilibrium scale.

For the radiation-reaction side of that story, the most appropriate benchmark is not the original Abraham–Lorentz equation but the reduced-order Landau–Lifshitz form, which avoids the familiar runaway and pre-acceleration pathologies of the AL/LAD hierarchy while preserving the leading radiation-reaction content [11,30]. In the nonrelativistic limit relevant for the present discussion one may write schematically

$$m\ddot{x} = F_{\text{ext}}(x, t) + \tau_0 \dot{F}_{\text{ext}}(x, t), \quad \tau_0 = \frac{q^2}{6\pi\epsilon_0 mc_0^3} \quad (16)$$

Conceptually, this is close to the Ford–O’Connell perspective, in which radiation reaction is written in a way that is naturally compatible with open-system and Langevin descriptions of a charge interacting with a bath [31]. That viewpoint is especially congenial here because my model already treats matter as an oscillator embedded in a stochastic environment. The point is not that radiation reaction alone proves gravity. The point is that it provides the right structural lesson: inertial response, dissipative response, and scale setting are all linked through the way matter exchanges power with its surroundings.

If the coarse-grained response of ordinary matter collapses to a common scalar control field $K(x)$ or $\zeta(x)$, then leading-order free fall is universal because the local acceleration is determined by the gradient of the equilibrium state rather than by a material-specific force coefficient. In the weak-field branch this may be summarized by

$$\mathbf{g}_{\text{eff}}(x) = -\nabla\Phi_\zeta(x), \quad \Phi_\zeta(x) \equiv -\frac{c_0^2}{2}\zeta(x)^2 \quad (17)$$

Free fall is then interpreted as a tiny contraction in the equilibrium scale of matter as it moves down a gradient in $\Phi\zeta$. What is ordinarily called gravitational potential energy is represented, in this reading, as a change in the internal equilibrium state of atoms and particles rather than as a localized substance of spacetime itself. The nontrivial requirement, of course, is universality: any material dependence hidden in the response kernels must coarse-grain away to within Eötvös-type experimental bounds [29]. That is why the equivalence principle is not assumed here as magic, but treated as a constraint that any successful microscopic completion must satisfy.

$$S_{\text{env}}(\omega, \mathbf{x}) = S_{\text{ZPF}}(\omega) + \delta S_{\text{matter}}(\omega, \mathbf{x}), \quad \zeta(\mathbf{x})^2 = \mathcal{F}[S_{\text{env}}] \quad (18)$$

Figure 3. Control hierarchy: ambient high-frequency bath versus low-frequency spectral weight




Figure 3. Ambient high-frequency bath versus low-frequency probe or control window.

6. Minimal propositions and proofs

Proposition 1. The operational scaling map is compatible with the Heisenberg products. *Proof.* Choose the length and time scalings from the adopted K-table. Then choose the complementary momentum and energy scalings so that the products remain unchanged. Direct multiplication gives Eqs. (3) and (4). No further assumptions are required. [14-17,21]

Proposition 2. The damping map reproduces the operational K-table if $K=(1-\zeta^2)^{-1}$. *Proof.* Substitute the defining relation for K into Eqs. (7)-(10). The frequency and energy rows immediately reproduce the $1/\sqrt{K}$ behavior, while the velocity, power, and acceleration rows reproduce the $1/K$ and $1/K^{3/2}$ behavior shown in Table 1. The effective coordinate mass parameter follows as the reciprocal power needed to preserve the force row. The linear-response definitions of Sec. 5.1 do not alter this algebraic matching; they only supply a more explicit interpretation of the control variable. [22,23]

Proposition 3. In the static spherical weak-field limit, the choice $\zeta^2(r)=2GM/(c^2r)$, or equivalently the closure of Sec. 5.2 with a point-mass source, reproduces the usual leading-order potential dependence of the K-table. *Proof.* Substitute Eq. (19) into Eq. (10) and expand for $2GM/(c^2r) \ll 1$. Equation (21) follows directly. The Earth-surface values quoted in Eq. (20) are then obtained by numerical substitution of $M \oplus$ and $R \oplus$. [3-5,22,23]

Proposition 4. If $K(x)$, or equivalently $\zeta(x)$, acts as a common scalar scaling field for all matter processes, then the leading-order free-fall response is composition independent. *Proof.* In the weak-field closure the acceleration field is $\mathbf{g}(\mathbf{x}) = -\nabla\Phi - \zeta(\mathbf{x})(c^2/2)\nabla\zeta^2(\mathbf{x})$, which contains no test-mass parameter. Composition dependence can therefore enter only through higher-order material response corrections hidden in the kernels defining $\zeta(x)$. The corresponding Eötvös parameter must then remain below existing bounds. [22,23,29]

These propositions do not complete a microscopic theory of gravity. What they establish is an internally coherent ladder of equivalences: metric scaling $\leftrightarrow$ K-scaling $\leftrightarrow$ uncertainty-compatible scaling $\leftrightarrow$ damping-compatible scaling. They also make plain where the program remains incomplete: the source law,

the detailed response kernels, the emergence of universality within experimental bounds, and the eventual requirement of a measurable departure from standard GR plus QED.

7. Thermodynamic context

My heat-bath analogy is not merely rhetorical. There is substantial literature showing that gravitational field equations can be regarded as thermodynamic or equation-of-state statements under suitable conditions. Jacobson's derivation of the Einstein equation from $\delta Q = TdS$ is the cleanest benchmark, and later entropic or emergent pictures of gravity extend the same broad lesson: macroscopic gravitational behavior may encode underlying microscopic degrees of freedom rather than exhaust them. [12,13]

This does not prove my model. It does, however, justify the style of explanation. If Einstein's equations can emerge as an equation of state, then it is reasonable to search for a deeper matter-and-environment description whose coarse-grained limit looks geometrical. My program belongs in that family of thought. Its distinctive claim is that the relevant coarse-graining may be written in terms of oscillator equilibrium, available driving power, and a real stochastic field environment that changes the scale of matter itself, while geometry serves as the macroscopic encoding of those changes. [12,13,22,23]

I also want that thermodynamic analogy stated carefully. I am not claiming that spacetime is literally a fluid or that thermodynamic language proves my microscopic picture. The point is narrower: if Einstein's equations can appear as an equation of state, then the metric may be a coarse-grained description of deeper degrees of freedom rather than the final ontology. That is the sense in which I use thermodynamics here. [12,13]

This also connects naturally to the vacuum-equilibrium strand of my own work. Puthoff's equilibrium description of radiationless ground states, together with the earlier harmonic-equilibrium picture in which particle scale is set by balance with a structured vacuum spectrum, suggests a common theme: matter properties may reflect a stable balance between internal dynamics and a surrounding field environment. My use of K and ζ can therefore be read as state variables of a matter-environment system, while geometry records the macroscopic comparison rules that emerge from it. [8,12,13,22-24]

Stated this way, the thermodynamic analogy motivates the present model without overclaiming. It tells me why an operational, matter-centered reinterpretation of GR is reasonable to pursue. It does not yet tell me the final microscopic field theory.

8. Weak-field matching and Earth example

For a static spherical source with negligible net charge, I adopt the identification $\zeta^2(r) = 2GM/(c^0 r)$. In that case $K(r) = 1/[1 - 2GM/(c^0 r)]$, and the operational relations reproduce the standard weak-field scalings of redshift and coordinate light speed in the chosen convention. At the surface of the Earth one obtains $\zeta \oplus \approx 3.73 \times 10^{-5}$ and a corresponding normalized frequency shift of order 10^{-9} , consistent with the fact that small fractional shifts can encode large macroscopic accelerations. [3-5,22,23]

This is an important engineering point. The model does not require large changes in dimensionless spectral quantities to produce the gravitational environment familiar in ordinary life. That observation, already present in my earlier papers, explains why the control problem is hard: even if the target shift is dimensionless and small per constituent, the total energy bookkeeping for macroscopic matter can still be enormous. [20,22,23]

The small-signal metrology relation follows directly from the scaling table. Since $v/v_0 = K^{(-1/2)} = \sqrt{1 - \zeta^2}$, a small engineered perturbation around $K \approx 1$ gives $\delta v/v \approx -(1/2)\delta K/K \approx -(1/2)\delta \zeta^2$. This is the practical equation to keep in mind when translating the model into a clock or spectroscopy experiment. [22,23,26-28]

This matters because modern optical clocks already operate in the 10^{-18} regime and have resolved relativistic redshift effects over laboratory-scale vertical separations. That does not validate my model by itself. It means only that the metrology is already sensitive enough to make a controlled search for a K-like perturbation a meaningful engineering question if such a perturbation can be generated cleanly. [26-28]

$$\zeta^2(r) = \frac{2GM}{c_0^2 r}, \quad K(r) = \frac{1}{1 - \frac{2GM}{c_0^2 r}} \quad (19)$$

$$z_\oplus \equiv 1 - \sqrt{1 - \zeta_\oplus^2} \approx 6.97 \times 10^{-10}, \quad \zeta_\oplus = \sqrt{\frac{2GM_\oplus}{c_0^2 R_\oplus}} \approx 3.73 \times 10^{-5} \quad (20)$$

$$K(r) \approx 1 + \frac{2GM}{c_0^2 r} + O(r^{-2}) \quad (21)$$

$$\nu/\nu_0 = K^{-1/2} = \sqrt{1 - \zeta^2}, \quad \delta\nu/\nu \approx -\frac{1}{2}\delta K/K \approx -\frac{1}{2}\delta\zeta^2 \quad (22)$$

9. Resonance, spectroscopy, and experimental program

The experimental language also benefits from sharpening. NMR and EPR are not Compton-frequency technologies, and I do not claim otherwise. The cleaner statement is that low-frequency laboratory probes may act as control interfaces into a much broader environmental coupling kernel. They can perturb state populations, coherence times, linewidths, relaxation channels, and possibly the effective damping parameter, without themselves being identified with the highest-frequency content of the ambient bath. [10,11,20,22,23]

That distinction suggests a realistic experimental program. I would not begin by claiming artificial gravity. I would begin by searching for anomalous, geometry-like frequency shifts or clock-comparison effects that remain after ordinary Zeeman, Stark, Lamb, cavity, thermal, and mechanical systematics have been removed. The targets are precision spectroscopy, resonators with tunable quality factor, atomic and solid-state clocks in engineered electromagnetic environments, and materials whose internal relaxation channels can be modulated reproducibly. Any positive claim must be differential, repeatable, and demonstrably larger than standard electromagnetic back-action. [10,11]

The resonance program is therefore a probe of coupling, not a proof of any active non-equilibrium engineering application. If an anomaly were found, it would still need to be mapped back into the K-table, checked for universality across materials, and tested against equivalence-principle bounds. [20,22,23]

The first discriminant is universality. A genuine K-like effect should shift distinct transitions by the same fractional amount when those transitions are exposed to the same engineered environment, because the hypothesis concerns the local scale of the clock sector itself. By contrast, ordinary Zeeman, Stark, and Lamb-type shifts carry species- and transition-dependent coefficients. [11,26-28]

The second discriminant is the geometry of comparison. Because a universal K-like perturbation rescales the local clock sector, two dissimilar clocks placed inside the same engineered region may preserve their ratio to leading order. The more decisive observable is comparison between a clock inside the engineered

environment and an external reference, or between two regions whose loading protocols differ in a controlled way. [26-28]

The third discriminant is protocol. Because the hypothesis concerns damping and loading, not merely static field amplitude, the most informative searches modulate cavity Q , linewidth, relaxation pathways, or spectral loading while conventional field amplitudes are held as fixed as possible. The signal to look for is a residual geometry-like shift that tracks the loading protocol rather than the ordinary electromagnetic coefficients alone. [10,11,20,22,23,25-28]

10. Scope of applicability and non-equilibrium extensions

The present paper establishes the passive branch of the model. With $0 \leq \zeta < 1$ and $K = (1 - \zeta^2)^{-1}$, the operational map implies $K \geq 1$. This is the branch that reproduces ordinary gravitational slowing of clocks, contraction of lengths, and the static/weak-field matching developed in the preceding sections. It is the branch for which the source law and damping identification have actually been written here. [3-5,20,22,23]

That scope matters. The closures adopted in this paper are phenomenological weak-field constructions, not yet a completed strong-field theory. Their purpose is to recover the observed static gravitational relations and to define a disciplined metrology program for spectroscopy, clock comparison, and resonance-based perturbation studies. The natural boundary of the present branch is $\zeta \rightarrow 1^-$, for which $K \rightarrow \infty$. How the model should be continued through horizons, or replaced by a more complete microscopic theory in strong fields, remains future work. [3-5,20,22,23]

For the same reason, I do not advance any $K < 1$ engineering claim in this paper. A driven or active non-equilibrium regime with $K < 1$ would require a different control law, a separate derivation, and a demonstration that the effect is not ordinary electromagnetic back-action dressed up in new language. The present paper therefore keeps its focus on the ordinary branch, the operational reinterpretation of GR within that branch, and the search for small anomalous K -like shifts in precision systems. [10,11,20,22,23]

11. Discussion

The virtue of the present synthesis is that it preserves the conceptual core of my work while stating it in cleaner and more rigorous terms. I am not claiming that all of quantum gravity has been solved. I am claiming that a non-geometrical interpretation of gravitational observables can be written coherently, that the uncertainty products can be preserved exactly under the adopted scaling, that the damping model reproduces the same transformation table once the identification $K = (1 - \zeta^2)^{-1}$ is made, and that the same weak/static observables of GR can therefore be read through a different ontology.

That point deserves to be said plainly. In the regime treated here, my model is not offered as a rival set of weak-field predictions to general relativity. It is a reinterpretation of the same successful relations. Wherever the same field equations and scaling laws are recovered, the physical predictions are identical. What differs is what those equations are taken to mean.

On this operational reading, Einstein's equation is not first treated as proof that a spacetime manifold is the fundamental substrate of reality. It is treated as a compact tensor map relating the observed relational structure of clocks, rulers, signals, and motion to the matter-energy content of the fields from which those standards of observation are themselves built. This does not diminish the mathematics of GR. It reorders the explanatory priority. [3-5,12,19-23]

While the geometrical interpretation has enriched the mathematics of gravitation enormously, its ontological primacy may also have constrained the search for quantum gravity by encouraging us to quantize geometry first and ask operational questions later. My claim is not that geometry is useless, but that it may be macroscopic bookkeeping rather than microscopic cause.

The burdens that remain are clear. First, the environmental field variable requires a proper source law. Second, the relation between $S_{\text{env}}(\omega, x)$ and the damping variable must ultimately be derived rather than phenomenologically weighted. Third, the universality of free fall must emerge or be explained within experimental bounds. Fourth, the model must predict a measurable deviation from standard GR plus QED if it is to move from interpretation to testable theory. Fifth, a full strong-field completion lies beyond the weak/static branch developed here. None of those are small tasks. But they are the right tasks, and stating them plainly is part of what turns a speculative idea into a research program.

12. Conclusions

The work assembled here shows that my program has a coherent and defensible core. The observed weak/static effects of gravitation may be written operationally through a single scalar variable K . The same table can be made fully compatible with the Heisenberg products by assigning the complementary momentum and energy scalings appropriately. The same table can then be matched again by a damped-oscillator description through the identification $K=(1-\zeta^2)^{-1}$. In that sense my long-standing claim survives careful scrutiny: there exists an engineering reinterpretation of gravitational scaling in which clocks, rulers, energies, and frequencies change because matter is operating at a different equilibrium scale, while geometry serves as the macroscopic encoding of those changes.

My claim, then, is not that general relativity has been overthrown, but that its equations admit a disciplined operational reading that is at least as natural for engineering purposes as the standard geometrical one. In the regime treated here, the model is intended to recover the same physical predictions as GR because it is a reinterpretation of those same relations. What changes is the ontology: the metric is read as a comparative map of physical clocks, rulers, signals, and energies rather than as direct evidence that a spacetime manifold must itself be the fundamental ontology.

This paper has also clarified what must be said more precisely than before. The controlling field is treated as a real effective stochastic background variable whose spectral and stochastic descriptions refer to the same underlying vacuum environment at different levels of description. Thermodynamic analogies motivate the search for an underlying layer, but do not prove it. NMR and EPR are best understood as control probes of coupling, not as direct access to Compton-scale carriers. Any active $K < 1$ extension lies outside the passive branch developed here and is not advanced in this paper. Even with those cautions, I regard the interpretive model as valuable. It places the emphasis where an engineer naturally wants it: on what clocks do, what rulers do, how matter is scaled, where the power flows, and which parameters one would have to control in order to move from observation toward technology. If a deeper non-geometrical account of gravitation is ever found, it will have to recover the same observational table. My claim is that the framework developed here is one disciplined way to begin that search.

Appendix A. Compact derivations

A.1 Heisenberg-product preservation

Let the observed coordinate comparison be length contraction by $1/\sqrt{K}$ and clock slowing by $\sqrt{K}$. Then choose the complementary uncertainty scalings $\Delta p_K = \sqrt{K} \Delta p_0$ and $\Delta E_K = \Delta E_0 / \sqrt{K}$. Direct multiplication gives

$$\begin{aligned}\Delta x_K \Delta p_K &= (\Delta x_0 / \sqrt{K}) (\sqrt{K} \Delta p_0) = \Delta x_0 \Delta p_0, \\ \Delta t_K \Delta E_K &= (\sqrt{K} \Delta t_0) (\Delta E_0 / \sqrt{K}) = \Delta t_0 \Delta E_0.\end{aligned}$$

Thus the adopted gravitational comparison table is compatible with the standard Heisenberg products without modifying the uncertainty principle itself.

A.2 Damping-to-K matching

The engineering model introduced here uses a phenomenological damping factor ζ for which the frequency and energy scales are multiplied by $\sqrt{1-\zeta^2}$ while time scales by its reciprocal. Defining $K = (1-\zeta^2)^{-1}$ gives immediately

$$\begin{aligned}\sqrt{1-\zeta^2} &= 1/\sqrt{K}, \\ 1/\sqrt{1-\zeta^2} &= \sqrt{K}, \\ (1-\zeta^2) &= 1/K, \\ (1-\zeta^2)^{3/2} &= 1/K^{3/2}.\end{aligned}$$

Every row of Table 1 then follows by direct substitution.

A.3 Weak-field spherical identification

For the static spherical matching used here, set $\zeta^2(r) = 2GM/(c_0^2 r)$. Then

$$K(r) = 1/(1-\zeta^2) = 1/[1 - 2GM/(c_0^2 r)].$$

Expanding for weak field, $2GM/(c_0^2 r) \ll 1$, gives $K \approx 1 + 2GM/(c_0^2 r) + O(r^{-2})$. This is the expected leading-order potential dependence in the chosen convention.

Appendix B. Phenomenological closures and experimental estimates

B.1 Weak-field source law and spherical reduction

The simplest closure that preserves the present interpretive program is to identify the damping variable with a weak-field potential Φ_ζ in the same way the Newtonian potential appears in the leading-order Schwarzschild / PV map. I therefore write

$$\zeta^2(\mathbf{x}) \equiv -2\Phi_\zeta(\mathbf{x})/c_0^2, \quad \nabla^2 \Phi_\zeta(\mathbf{x}) = 4\pi G \rho_{\text{eff}}(\mathbf{x}) \quad (\text{B1})$$

and take the effective source density to include both ordinary matter and the matter-induced departure of the local field environment from the baseline vacuum state:

$$\begin{aligned}\rho_{\text{eff}}(\mathbf{x}) &= \rho_m(\mathbf{x}) + \lambda \delta u_{\text{env}}(\mathbf{x})/c_0^2, \\ \delta u_{\text{env}}(\mathbf{x}) &= \int_0^\infty W_g(\omega) \delta S_{\text{matter}}(\omega, \mathbf{x}) d\omega\end{aligned} \quad (\text{B2})$$

For a static spherical source with $\rho_{\text{eff}}(\mathbf{x}) = M \delta^3(\mathbf{x})$, the usual potential is recovered and the matching used in the main text follows immediately:

$$\Phi_\zeta(r) = -GM/r, \quad \zeta^2(r) = 2GM/(c_0^2 r), \quad K(r) = 1/[1 - 2GM/(c_0^2 r)] \quad (\text{B3})$$

This appendix does not pretend that the microscopic source law is finished. It makes explicit the minimal closure already implicit in the weak-field matching of the main paper. [3-5,20,22-25]

B.2 Linear-response bridge from $S_{\text{env}}(\omega, \mathbf{x})$ to $\zeta(\mathbf{x})$

The environmental field variable is most naturally separated into a spectral state variable and a response variable. A coarse-grained local environmental energy density may be defined by

$$u_{\text{env}}(\mathbf{x}) = \int_0^\infty W(\omega) S_{\text{env}}(\omega, \mathbf{x}) d\omega \quad (\text{B4})$$

while the effective damping rate and damping ratio are written phenomenologically as

$$\begin{aligned} \gamma_{\text{eff}}(\mathbf{x}) &= \gamma_0 + \int_0^\infty G_\gamma(\omega) \delta S_{\text{env}}(\omega, \mathbf{x}) d\omega, \\ \zeta(\mathbf{x}) &= \gamma_{\text{eff}}(\mathbf{x})/\omega_0 \end{aligned} \quad (\text{B5})$$

The corresponding local response function of a bound mode then takes the usual damped-oscillator form

$$\chi(\omega; \mathbf{x}) = 1/\{m[\omega_0^2 - \omega^2 - i2\gamma_{\text{eff}}(\mathbf{x})\omega]\} \quad (\text{B6})$$

which shows explicitly how the spectral environment and the damping parameter enter at different conceptual levels. The environment is the bath; ζ is the scalar summary of the bath's effect on the mode of interest. [8-10,20,22,23,25]

B.3 Universality and the Eötvös parameter

If the K-map is common to all matter processes, then leading-order free fall is universal. Composition dependence may be parameterized by small residual response coefficients ε_a and ε_b :

$$\begin{aligned} g_a &= g(1 + \varepsilon_a), \quad g_b = g(1 + \varepsilon_b), \\ \eta_{ab} &= 2|g_a - g_b|/|g_a + g_b| \approx |\varepsilon_a - \varepsilon_b| \end{aligned} \quad (\text{B7})$$

In this framework the burden is therefore clear: any material dependence hidden in the response kernels must be suppressed sufficiently that η_{ab} remains below present experimental limits. The final MICROSCOPE result gives the relevant order of magnitude for that requirement. [29]

B.4 Small-signal spectroscopic estimate

For a small engineered perturbation about $K \approx 1$, the metrology equation is immediate from the scaling map:

$$\nu/\nu_0 = K^{-1/2} = \sqrt{1 - \zeta^2}, \quad \delta\nu/\nu \approx -\frac{1}{2} \delta K/K \approx -\frac{1}{2} \delta\zeta^2 \quad (\text{B8})$$

This is the direct bridge between the interpretive model and clock-based experiments. Modern optical clocks and clock comparisons are already sensitive enough to make such a search technically meaningful, provided conventional electromagnetic systematics are controlled to the same level. [26-28]

B.5 Strong-field outlook: the critical-damping boundary

The passive oscillatory branch developed in this paper is defined by $0 \leq \zeta < 1$ and by the real-frequency identification $\omega_\zeta = \omega_0 \sqrt{1 - \zeta^2}$. If the same spherical damping map is formally continued beyond the weak-field regime, then the model supplies a natural strong-field boundary of its own. Define the Schwarzschild radius in the usual way and rewrite the spherical matching in that language. [3-5,22,23]

$$r_s = \frac{2GM}{c_0^2} \quad (\text{B9})$$

$$\zeta^2(r) = \frac{2GM}{c_0^2 r} = \frac{r_s}{r} \quad (\text{B10})$$

$$K(r) = \frac{1}{1 - r_s/r} \quad (\text{B11})$$

The horizon is then the point at which the underdamped branch reaches $\zeta=1$. In the present comparison scheme this is also the point at which the external static scaling parameter diverges. Approached from the exterior static branch, the limit from above implies $K \rightarrow \infty$.

$$r = r_s \Rightarrow \zeta = 1 \quad (\text{B12})$$

$$r < r_s \Rightarrow \zeta > 1 \quad (\text{B13})$$

In damping language this is a critical point. For radii below the Schwarzschild radius the continuation lies on the overdamped branch. Writing the unforced part of Eq. (6) in characteristic form gives

$$\lambda^2 + 2\zeta\omega_0\lambda + \omega_0^2 = 0 \quad (\text{B14})$$

$$\lambda_{\pm} = -\omega_0(\zeta \pm \sqrt{\zeta^2 - 1}) \quad (\text{B15})$$

Thus the ordinary real-frequency map used in the main text terminates at the critical point and must be replaced, inside the formal continuation, by a relaxation-rate description. I do not present this appendix as a completed interior black-hole solution, nor do I claim that a freely falling observer encounters a local singular dissipation at the horizon. The point is narrower: the same damping model used for the passive gravitational branch contains a natural strong-field boundary at $\zeta=1$, with an overdamped continuation below the Schwarzschild radius. That observation suggests a path for extending the model beyond the strict weak-field regime without altering its operational starting point. [3-5,22,23]

References

- [1] Wilson, H. A. An electromagnetic theory of gravitation. *Physical Review* 17, 54-59 (1921).
- [2] Dicke, R. H. Gravitation without a principle of equivalence. *Reviews of Modern Physics* 29, 363-376 (1957).
- [3] Einstein, A. On the influence of gravitation on the propagation of light. *Annalen der Physik* 35, 898-908 (1911).
- [4] Puthoff, H. E. Polarizable-vacuum (PV) representation of general relativity. *arXiv:gr-qc/9909037* (1999).
- [5] Puthoff, H. E. Polarizable-vacuum (PV) approach to general relativity. *Foundations of Physics* 32, 927-943 (2002).
- [6] Puthoff, H. E., Maccone, C. & Davis, E. W. Levi-Civita effect in the polarizable-vacuum representation of general relativity. *General Relativity and Gravitation* 37, 483-489 (2005).
- [7] Puthoff, H. E., et al. Engineering the zero-point field and polarizable vacuum for interstellar flight. *Journal of the British Interplanetary Society* 55, 137-144 (2002).
- [8] Puthoff, H. E. Quantum ground states as equilibrium particle-vacuum interaction states. *Quantum Studies: Mathematics and Foundations* 3, 5-10 (2016).
- [9] Milonni, P. W. Quantum mechanics of the Einstein-Hopf model. *American Journal of Physics* 49, 177-181 (1981).
- [10] Milonni, P. W. *The Quantum Vacuum: An Introduction to Quantum Electrodynamics*. Academic Press (1994).
- [11] Jackson, J. D. *Classical Electrodynamics*, 3rd ed. Wiley (1999).
- [12] Jacobson, T. Thermodynamics of spacetime: the Einstein equation of state. *Physical Review Letters* 75, 1260-1263 (1995).

- [13] Verlinde, E. On the origin of gravity and the laws of Newton. *Journal of High Energy Physics* 2011, 029 (2011).
- [14] Adler, R. J. & Santiago, D. I. On gravity and the uncertainty principle. *Modern Physics Letters A* 14, 1371-1381 (1999).
- [15] Kuzmichev, V. E. & Kuzmichev, V. V. Uncertainty principle in quantum mechanics with Newton's gravity. *European Physical Journal C* 80, 248 (2020).
- [16] McCulloch, M. E. Gravity from the uncertainty principle. *Astrophysics and Space Science* 349, 957-959 (2014).
- [17] McCulloch, M. E. Quantised inertia from relativity and the uncertainty principle. *EPL* 115, 69001 (2016).
- [18] Alcubierre, M. The warp drive: hyper-fast travel within general relativity. *Classical and Quantum Gravity* 11, L73-L77 (1994).
- [19] Desiato, T. J. General Relativity and the Polarizable Vacuum. Manuscript (2006).
- [20] Desiato, T. J. The Electromagnetic Quantum Vacuum Warp Drive. *Journal of the British Interplanetary Society* 68, 347-353 (2016).
- [21] Desiato, T. J. The Derivation of Gravity from the Uncertainty Principle: Introducing the Maxwell Temporal Field. Manuscript v12 (2021).
- [22] Desiato, T. J. Engineering a Warp Drive Using Quantum Gravity and a New Interpretation of General Relativity: An Engineering Model. Manuscript updated 7 Oct 2023.
- [23] Desiato, T. J. An Engineering Model of Quantum Gravity. Manuscript updated 18 Sept 2016.
- [24] Desiato, T. J. & Storti, R. C. Electro-Gravi-Magnetics (EGM): The Harmonic Representation of Particles. Manuscript v1 (2005).
- [25] Callen, H. B. & Welton, T. A. Irreversibility and generalized noise. *Physical Review* 83, 34-40 (1951).
- [26] Ludlow, A. D., Boyd, M. M., Ye, J., Peik, E. & Schmidt, P. O. Optical atomic clocks. *Reviews of Modern Physics* 87, 637-701 (2015).
- [27] Chou, C. W., Hume, D. B., Rosenband, T. & Wineland, D. J. Optical clocks and relativity. *Science* 329, 1630-1633 (2010).
- [28] McGrew, W. F. et al. Atomic clock performance enabling geodesy below the centimetre level. *Nature* 564, 87-90 (2018).
- [29] Touboul, P. et al. MICROSCOPE Mission: Final Results of the Test of the Equivalence Principle. *Physical Review Letters* 129, 121102 (2022).
- [30] Landau, L. D. & Lifshitz, E. M. *The Classical Theory of Fields*, 4th ed. Butterworth-Heinemann (1980).
- [31] Ford, G. W. & O'Connell, R. F. Radiation reaction in electrodynamics and the elimination of runaway solutions. *Physics Letters A* 157, 217-220 (1991).